

Through Thick and Thin

Learning to Live with Warfarin

South Denver Cardiology Associates PC

What is Anticoagulation?

- Anti = against
- Coagulation = blood clotting
- An anticoagulant is a medication that reduces the blood's potential to clot
- Anticoagulants are often called "blood-thinners"

Reasons for Anticoagulation Therapy

- Deep Vein Thrombosis (DVT)
- Mechanical Heart Valve Replacement
- Pulmonary Embolus (PE)
- Atrial Fibrillation (Afib)
- Atrial Flutter (Aflutter)
- Myocardial Infarction (MI) or Heart Attack

3

Reasons for Anticoagulation Therapy

- Cardiomyopathy or Left Ventricular Dysfunction
- Stroke
- Transient Ischemic Attack (TIA)
- Post-operative

4

About Warfarin (Coumadin®)

- Most commonly used oral anticoagulant
- Stops harmful blood clots from forming
- Partially blocks the re-use of vitamin K in the liver
 - vitamin K is needed to help the blood form clots to prevent bleeding
 - vitamin K is found in green leafy vegetables

5

About Warfarin (Coumadin®)

- Warfarin tablets
 - numerical imprint of strength
 - 1mg, 2mg, 2.5mg, 3mg, 4mg, 5mg, 6mg, 7.5mg, or 10mg
 - color-coded
 - scored in the middle
- Important to pay attention to any change in size, color or shape of tablets
 - can affect therapeutic dose

6

About Warfarin (Coumadin®)

- Warfarin therapy monitored by blood test called INR (International Normalized Ratio)
 - MD will prescribe what range INR needs to be
 - INR = 0.9-1.1, without warfarin
 - INR = 2.0-3.0, for most indications
 - INR = 2.5-3.5, for mechanical heart valves and other conditions
 - general guidelines, may have different range according to medical condition or other risk factors

7

About Warfarin (Coumadin®)

- Warfarin has a “narrow therapeutic range”
- When INR falls within prescribed range, level is “therapeutic”
- When INR falls below prescribed range:
 - INR = 1.5, for example
 - blood is “too thick”, clotting too fast
 - risk for forming blood clots exists
 - warfarin dose needs to be increased

8

About Warfarin (Coumadin®)

- When INR is above prescribed range:
 - INR = 4.5, for example
 - blood is “too thin”, clotting too slowly
 - higher risk for bleeding
 - warfarin dose needs to be decreased

9

About Warfarin (Coumadin®)

- The dose of warfarin needed is the one that will keep INR in therapeutic range
 - some people need very large dose of warfarin
 - some people need very small dose of warfarin
- The dose of warfarin will fluctuate with changing INR values
 - dose likely to change with each INR in first month of therapy
 - will change after first month with changes in diet, medications, activity and overall health

10

About Warfarin (Coumadin®)

- IT IS VERY IMPORTANT TO TAKE WARFARIN AS PRESCRIBED!
 - Make sure you understand exactly how much warfarin you need to take every day until next INR is scheduled
- It is also important to take warfarin at the same time every day
 - preferably in the evening

11

About Warfarin (Coumadin®)

- The INR will need to be checked
 - 2-3 times in the first week of warfarin therapy
 - to once a week until stable (usually about a month)
- If INR becomes stable quickly, INR needs to be checked less often
- If INR does not become stable, INR will need to be checked more frequently

12

About Warfarin (Coumadin®)

- When a stable dose of warfarin has been found
 - INR needs to be checked once a month
 - If INR is stable for several consecutive months a 6-week schedule may be considered for INR's
- It is very important to call for earlier INR appointment if there are changes
 - in diet, medications, activity and overall health

13

About Warfarin (Coumadin®)

- If a dose of warfarin is missed it can be taken later the same day
- If previous days' dose was missed
 - DO NOT TAKE A DOUBLE DOSE!
 - Get back on dosing schedule and let healthcare provider know the date of the missed dose with next INR

14

About Warfarin (Coumadin®)

- If more than 1 or 2 doses are missed
 - call healthcare provider to discuss
 - INR may need to be checked
- You are not likely to have symptoms if INR is too low or too high
- This is why regular testing of INR is so important

15

Warfarin and Side Effects

- The most common and serious side effect of warfarin therapy is bleeding
 - The risk of bleeding is relatively small when warfarin is taken properly
 - To lower the risk, make sure to have INR levels checked regularly
 - Report any changes in diet, medications, activity and overall health

16

Warfarin and Side Effects

- Monitor and report any of the following
 - Prolonged or unusual bleeding from cuts
 - Nosebleeds
 - Bleeding from gums when brushing teeth
 - Increased menstrual flow or unexpected vaginal bleeding
 - Bright red, tea-colored, or dark brown urine
 - Bright red or black, tarry stools

17

Warfarin and Side Effects

- Monitor and report any of the following
 - Prolonged severe and unusual headache or stomachache
 - Unusual weakness, dizziness or changes in vision or speech
 - Coughing or spitting up blood
 - Vomiting bright red or coffee-ground emesis
 - Joint pain or immobility
 - Unusual pain or swelling in extremities

18

Warfarin and Side Effects

- Any of these symptoms could be significant, but may not be dangerous
- In therapeutic doses warfarin does not cause bleeding, but will make it harder to stop bleeding if an injury occurs
- Primary care physician (PCP) will need to be aware of, and treat any of these side effects

19

Warfarin and Side Effects

- For minor bleeding
 - nosebleed or bleeding gums that stop within a few minutes
- Call healthcare provider so INR can be checked
- Reporting bleeding symptoms early and having warfarin dose adjusted
 - can prevent INR from increasing and more serious side effects from occurring

20

Warfarin and Side Effects

- GET TO THE NEAREST EMERGENCY ROOM
 - If you have major bleeding
 - Nosebleed that does not stop after 15-20 minutes
 - Frank blood in stools, or black, tarry stools
 - Vomiting bright red or coffee-ground emesis
 - If you sustain a traumatic injury from a fall or car accident
 - You will need to be checked for internal bleeding

21

Warfarin and Side Effects

- GET TO THE NEAREST EMERGENCY ROOM
 - If you are exhibiting signs of a stroke or “brain attack”
 - Sudden numbness or weakness of face, arm or leg
 - Difficulty speaking or understanding others
 - Problems seeing clearly
 - Confusion, dizziness, severe drowsiness
 - Loss of coordination
 - Sudden severe headache with no known cause

22

Warfarin and Side Effects

- Rash
 - Please report if develop rash starting warfarin
 - Could be minor
 - Could be allergy to dye in pills

23

Special Considerations with Warfarin

- Pregnancy
 - Report any plans to become pregnant or pregnancies immediately
 - Warfarin is very harmful to the developing fetus
- Avoid activities that put you at risk for traumatic injury
- Exercise caution when using sharp objects

24

Special Considerations with Warfarin

- Advisable to wear a Medic-Alert bracelet
 - To notify emergency healthcare personnel you are taking anticoagulants
- Have an updated medication list
 - In wallet or billfold and with you at all times
- Be careful! Use common sense!
 - You are taking a blood thinner and are going to bruise more easily and bleed longer than before you took warfarin

25

Warfarin and Drug Interactions

- Warfarin has hundreds of drug interactions
 - With prescription, over-the-counter and herbal medicines
 - Not all medications affect everyone the same
 - Initiation of some meds warrant an INR within 2-3 days
 - Initiation or change in others may need INR to be checked within 2 weeks

26

Warfarin and Drug Interactions

- One of the most preventable causes of an untherapeutic INR
 - Is to inform healthcare provider when making any medication changes, however minor they may seem
- If you need to take a medication that interacts with warfarin
 - You will need more frequent testing of INR and adjustment in dose until INR is stable

27

Warfarin and Diet

- Warfarin works against vitamin K in diet
 - Vitamin K is important in forming blood clots
 - Foods with the highest amounts of vitamin K are green, leafy vegetables
- Green vegetables are important!
 - Contain folic acid, phytonutrients and fiber
 - All of which help protect your heart and have countless other benefits to body
 - Eat your veggies!

28

Warfarin and Diet

- The most important thing is to maintain a consistent intake of vitamin K in diet
 - Eat same quantity and quality of green vegetables weekly
 - If you eat more green veggies than usual
 - INR likely to drop, putting you at risk for clots
 - If you eat less green veggies than usual
 - INR can increase, putting you at risk for bleeding
 - If you eat the same amount of green veggies
 - It will help to balance warfarin dose

29

Warfarin and Diet

- Not all green vegetables have the same amount of vitamin K
 - As a general rule, the greener and leafier it is, the more vitamin K it contains
 - Green beans and carrots are “low” in vitamin K
 - Iceberg lettuce and asparagus are ‘medium’ in vitamin K
 - Spinach and broccoli are ‘high’ in vitamin K
- Vitamin K is also found in
 - Oils, olive and soybean are ‘high’
 - Margarine and mayonnaise are also ‘high’

30

Warfarin and Dietary Supplements

- Dietary supplements such as Ensure, Boost, Carnation Instant Breakfast, and various protein drinks are 'high' in vitamin K
 - If you start a dietary regimen with any of these
 - Be consistent
 - Call healthcare provider, as warfarin dose will likely need to be adjusted

31

Warfarin and Vitamins

- Multi-vitamins (MVI)
 - Contain vitamin K
 - Call healthcare provider if starting or changing
 - Pay attention to vitamin K content when choosing or switching brands
 - Viactiv MVI and Calcium chews have extremely high amounts of vitamin K
- B Vitamins
 - Have little to no effect on warfarin

32

Warfarin and Vitamins

- Vitamin C
 - Up to 500mg daily has no affect, greater than 500mg daily may effect warfarin dosage
- Vitamin E
 - Up to 400 iu daily has no affect, doses greater than that may increase INR
- Fish oil
 - Likely to increase INR and require dose adjustment

33

Warfarin and Alcohol

- Drinking alcohol can affect INR and is not always predictable
- Best not to drink alcohol, but alcoholic beverages in moderation can be safe
- One or two drinks each day
 - 12 oz. Beer = 4 oz. Wine = 1 oz. Liquor
- It is best to be consistent from week to week

34

Warfarin and Alcohol

- Excessive alcohol intake
 - puts you at significant risk for injury and potential bleeding complications while taking warfarin
- Please call to have INR checked
 - If you make any changes in alcohol intake
 - Either an increase or decrease can affect INR and warfarin dose may need to be adjusted

35

Warfarin and Exercise

- Physical activity can affect INR levels
 - A change, either increase or decrease may require an adjustment in warfarin dose
- Please contact healthcare provider when making any significant changes in physical activity or exercise program

36

Warfarin and Medical, Surgical or Dental Procedures

- Contact your healthcare provider any time you need to have a procedure that puts you at risk for bleeding
 - Warfarin may need to be temporarily stopped
 - It may not be safe for you to stop your anticoagulants entirely
 - You may need an alternative method of anticoagulation

37

Warfarin and Travel

- Travel can often affect INR levels
- Have INR checked before travelling
- If an INR needs to be checked while away, or you will be gone for an extended time
 - Your healthcare provider can help you make arrangements to have it checked while away
- Try to maintain a consistent diet and avoid excessive alcohol intake

38



Questions?

39